COMPOSITE SECTION AND FAUNAL DISTRIBUTION CHART

LOWER ORDOVICIAN | MID. ORDOVICIAN

JEFFERSON CITY GROUP | BIG BUFFALO SERIES

R'd'x. Form. | Rich Fountain Formation 0'—180' | Theodosia Formation 0'—30' | Cotter Formation 0'—340' | Powell Formation 0'—214' | Everton Formation

LUTIE MEMBER 0'—170'

BLACKJACK KNOB MEMBER 0'—180'

Scale 50' 25' 0'

School Mine Ledge

Rockaway Cgl. beds

Hercules Tower Ss.

Pocket Hollow Oolite

Gainesville Ss.

Crooked Creek Chert

Jenkins Branch Chert

Bull Cryp Bed

Bull Shoals Mbr. Chert

Black Ledge

Ozarkocoelia
O. irregularis
Archaeoscyphia
A. annulata
Diporelasma
D. silicum
Tylocechia
T. typica
T. delicatula
Polytoechia
P. inaequistriata
Syntrophopsis
S. arkansasensis
S. grandis
Proplina
P. elongata
P. grandis
Raphistomina
R. inflata
Ophileta
O. sculpta
Orospira
O. blairanosa
O. gainesvillensis
O. depressa
O. elegantula
O. elevata
O. conica
Palhemia
P. taneyensis
Lesueurilla
L. grandis
L. isabellaensis
L. magna
L. planidorsalis
L. supraconvexa
Eotomaria
E. depressa
Hormotoma
H. cotterensis
H. dubia
H. lutiensis
H. powellensis
Geratopsa
G. ankylosa
G. calceoliformis
G. capuliformis
G. inflata
G. jonesi
G. subconica
G. supraplana
G. tennesseensis
G. torta
Shumardoceras
S. complanata
S. lutiensis
Aphetoceras
A. ozarkensis
A. subcostulatum
Campbelloceras
C. simplex
Centrotarphyceras
C. (?) baxterensis
C. yellvillensis
Pilotoceras
P. burnei
Moreauoceras
M. millerl
Descritoceras
D. jeffersonensis
Cyptendoceras
C. magnus
Bathyurellus
B. permarginatus
Jeffersonia
J. crassimarginata
J. delicatula
J. granata
J. jennii
J. mediorista
J. missouriensis
J. producta
J. sp. 1.
J. sp. 2.
Bathycephalus
B. jeffersonensis
B. st. clairi
Lutesvillia
L. bispinosa
Hystricurus
H. abruptus
H. conicus
Rollia
R. goodwini
Rananasus
R. brevicephalus
R. conicus
Genioteus
G. marionensis
G. semicircularis
gen. et sp. indet.

PLATE II